

**DIVERSITY OF THE *FUSARIUM VERTICILLIOIDES* AND
F. PROLIFERATUM ISOLATES ACCORDING TO THEIR FUMONISIN
B₁ PRODUCTION POTENCIAL AND ORIGIN**

Sonja TANČIĆ¹, Slavica STANKOVIĆ², Jelena LEVIĆ², Vesna KRNJAJA³,
Jelena VUKOJEVIĆ⁴

¹Laboratory of Phytopathology, Oilcrops Department, Institute of field and vegetable
crops, Novi Sad, Serbia

²Laboratory of Phytopathology, Maize Research Institute Zemun Polje, Serbia

³Institute of Animal Husbandry, Zemun, Serbia

⁴Department of Algaeology, Mycology and Lichenology, University of Biology
Belgrade, Belgrade, Serbia

Tančić S., S. Stanković, J. Lević, V. Krnjaja and J. Vukojević
(2012): *Diversity of the Fusarium verticillioides and F. proliferatum*
isolates according to their fumonisin B₁ production potencial and origin
- Genetika, Vol 44, No. 1,163 -176.

Species of the genus *Fusarium* are characterised by the exceptional
intraspecies and interspecies variability in respect to morphological,
physiological and genetic properties. Intraspecies and interspecies
diversity of *Fusarium verticillioides* and *Fusarium proliferatum* isolates
in the production of fumonisin B₁ according to their origin from maize
and wheat grains was studied. Fumonisin B₁ production potential of

investigated 42 isolates was assessed by HPTLC and ELISA method. All 22 and 20 investigated *F. verticillioides* and *F. proliferatum* isolates, respectively, had the ability to produce fumonisin B₁ toxin. Fumonisin B₁ production potential of *F. verticillioides* and *F. proliferatum* varied from 88.60 ppm to 1,300.60 ppm and from 2.37 ppm to 1,246.00 ppm, respectively. According to Mann-Whitnev U test, there were no significant differences between the fumonisin B₁ production mean values of isolates of *F. verticillioides* originated from maize and wheat (628.13 ppm and 696.38 ppm, respectively), as well as, fumonisin B₁ production mean values of *F. proliferatum* isolates (212.32 ppm and 158.07 ppm, respectively). Variability coefficient values indicated that fumonisin B₁ production potential variability was more expressed in maize than wheat originated isolates for both species.

Key words: *Fusarium verticillioides*, *F. proliferatum*, fumonisin B₁ maize, production variability, wheat

INTRODUCTION

Species of the genus *Fusarium* are characterised by the exceptional intraspecies and interspecies variability in respect to morphological, physiological and genetic properties. Due to their great variability, species from the section *Liseola* (*Fusarium verticillioides* Sheldon – syn. *F. moniliforme* Sheld., *F. subglutinans* (Wollen. and Reink.) Nelson, Toussoun and Marasas, *F. proliferatum* (Matsushima Nirenberg) are well known as pathogens of a great number of agricultural host plants such as maize (LOGRIECO *et al*, 2002; FANDOHAN *et al*, 2003), wheat (BIRZELE *et al*, 2002; LOGRIECO *et al*, 2003; FURLONG *et al*, 2005; KRNIJAJA *et al*, 2008), barley (BATATINHA *et al*, 2007; STANKOVIĆ *et al*, 2011), oat (BOTTALICO and PERRONE, 2002), hops (STANKOVIĆ *et al*, 2008a), sorghum (DA SILVA *et al*, 2006), asparagus (STĘPIEŃ *et al*, 2011), rice (DESJARDINS *et al*, 2000), bananas (MARASAS *et al*, 1998), mango (BRITZ *et al*, 2002), pineapple (STĘPIEŃ *et al*, 2011), etc. Although they are cosmopolites, some species are more frequent in certain climatic regions and certain host plants. They cause root, stem and ear rot with reductions in crop yields estimated between 10% and 30% in Europe (LOGRIECO *et al*, 2002).

Additionally to their economic importance, these species are capable to produce mycotoxins in pre-harvest infected plants or in stored grains. Fumonisin are a group of 15 structurally related mycotoxins, and three of them (FB₁, FB₂ and FB₃) can cause a serious health problem in animals such as equine leukoencephalomalacia (ELEM), pulmonary edema in swine, and liver or kidney cancer in rats or mice. Fumonisin are also associated with human esophageal cancer (DESJARDINS, 2006).

The FB₁ concentration depends on the origin of isolates, including a host plant and a geographic region. Under laboratory conditions, isolates of *F. verticillioides* and *F. proliferatum*, derived from maize, synthesise more FB₁ (average 1259 ppm) than the isolates originating from grains of wheat (average 769 ppm) or barley (average 320 ppm) (VISCONTI *et al*, 1994). NELSON *et al*. (1991) tested strains of *F. verticillioides* from various substrates and geographic areas for FB₁ production and

found that the most strains from maize-based feed (16/20) associated with leukoencephalomalacia were high producers, whereas among the strains isolated from millet and sorghum grain, were few producers (4/15). The strains from maize from Nepal had one high producer (1/10), and the most strains from maize of good quality intended for feed of poultry (8/9) and several strains from human tissue (9/13) were mainly high producers. Isolates from sorghum grain and trees, maize, sugar cane and from the Australian soil did not produce fumonisins (0/10).

Fungi produce fumonisin B₁ in the highest concentrations when grain moisture ranges from 27.0 to 32.0% (BARS *et al.*, 1994). High concentrations of fumonisin B₁ (10.242 ppm) and B₂ (3.068 ppm) were obtained when *F. verticillioides* had been cultivated on the maize substrate, rice medium (0.206 ppm FB₁, 0.100 ppm FB₂, respectively), and on the peanut and soybean substrate (\leq 0.05 ppm) (HOLCOMB *et al.*, 1993).

The most frequent species in Serbia identified on maize kernels were *F. graminearum* Schwabe, *F. verticillioides* and *F. subglutinans*, while on wheat kernels were *F. graminearum*, *F. poae* (Peck) Wollen. and *F. proliferatum* (LEVIĆ *et al.*, 2001; BOČAROV-STANČIĆ *et al.*, 2000). Although the frequency of FB₁ producers were determined on maize and wheat kernels in Serbia, a few studies had been carried out with the aim to observe their diversity in the FB₁ production (ĐILAS *et al.*, 2003; STANKOVIĆ *et al.*, 2008c; STANKOVIĆ *et al.*, 2008b).

The aim of our research was to determine genetic FB₁ production potential under the laboratory conditions of 42 *F. proliferatum* and *F. verticillioides* isolates in relation to their origin. This paper reports the intra- and interspecies variability in FB₁ production potential of *F. proliferatum* and *F. verticillioides* isolates originated from maize and wheat kernels collected in 10 different locations in Serbia during the period 2005-2007.

MATERIALS AND METHODS

Plant host samples

With the aim to obtain *F. verticillioides* and *F. proliferatum* isolates for this study, samples of maize and wheat kernels were collected during the 2005, 2006 and 2007 from Northern (Bačka Topola, Kikinda, Kovin, Novi Sad, Sombor, Sremska Mitrovica) and Central/Southern Serbia (Kraljevo, Loznica, Niš, and Šabac).

Wheat samples were based on a few smaller samples collected from different parts of stored kernels one month after the harvest in 2005 and 2006. Thirty two kernels from each subsample (four subsamples per location) were incubated at water agar (WA) in order to analyse diversity of *Fusarium* species.

Ten maize ears with *Fusarium* symptoms, collected diagonally from each field in 22 sites of 10 locations during the 2006 and 2007, were taken as a basic sample. Ten kernels were taken as subsamples from each ear (100 kernels per sample) and they were incubated at potato dextrose agar (PDA) for diversity analyses of *Fusarium* species.

Fungal isolates

For the fungal identification, 128 wheat kernels (32 per replication) and 100 maize kernels (25 per replication) of each sample were incubated for seven days on water agar (WA) and potato dextrose agar (PDA), respectively. *Fusarium* species were identified according to LESLIE and SUMMERELL (2006). All isolated *Fusarium* cultures were refined to monosporous isolates, and 22 isolates of *F. verticillioides* and 20 isolates of *F. proliferatum* were selected for further toxicological investigations analyses.

Sample preparation for fumonisin assessment

Fifty grams of maize kernels were soaked in sterile distilled water to reach 40% of moisture. After 24 hours, maize kernels were autoclaved for 30 min. Sterilised maize kernels were inoculated with 3-5 plugs (5x5 mm) of monosporous *F. verticillioides* or *F. proliferatum* isolates previously cultivated for seven days on PDA at 25°C in the dark. Inoculated flasks were incubated for 2-3 weeks at 27°C in the dark, occasionally shaken to prevent sample conglomeration and unequal mycelial growth, which could reduce fumonisin production. After incubation, infected maize kernels were dried at 50°C for 3 days and milled to powder, which was kept in plastic bags at 4°C till further fumonisin extraction.

Toxicology assessment

The FB₁ production potential of tested isolates was assessed by two methods: High Performance Thin Layer Chromatography (HPTLC) according to LOGRIECO et al. (1995) and Enzyme-linked immunosorbent assay (ELISA). The HPTLC method had three phases: preparation of samples, fumonisin extraction and purification, and HPTLC fumonisin detection. For fumonisin extraction, 70% methanol was used. Powdered samples and methanol (1:10) were blended for 3 min (1300 rpm) and filtered through Whatman filter paper No 1. Filtrates (10 ml) were purified by Bonded Phase Sax Columns and appropriate solvents in Solid-Phase Extraction Manifold. Tubes with extracted fumonisin were then evaporated in the vacuum evaporator at 50°C till the appearance of a dry residue at the bottom of the tube. The fumonisin dry residue was resolved in 1 ml of methanol, and 5 µl of a sample was used for the further fumonisin detection. HPTLC plates 20x10 cm Kiesgel 60 F254 MERC with the appropriate solvent, development solution and FB₁ standard, were used for fumonisin detection in samples.

The ELISA method also encompassed three phases: sample preparation, ELISA assay procedure and ELISA fumonisin detection. The ELISA assay procedure was performed according to CELER FUMO (code MF101) Test Kit Manual, while the ELISA fumonisin detection was done at 450 nm (Labsystems MultiScan® MCC/340).

Statistical analyses

Basic statistic analyses (mean values, variability coefficient, correlation between HPTLC and ELISA data) were done using software Statistica 7.0 (StatSoft)

and significant differences in mean values of the FB₁ production were ascertained by the Mann-Whitney U test.

RESULTS AND DISCUSSION

Due to the application of two different methods for the quantitative analysis of FB₁ (HPTLC and ELISA), the positive correlation between these methods ($p < 0.05$) was established by the statistical analysis. Coefficients of correlation between the HPTLC and ELISA were $r = 0.9633$ and $r = 0.9697$ for FB₁ production by *F. verticillioides* and *F. proliferatum*, respectively. The application of the ELISA method provided a greater reliability and preciseness, and the data obtained by ELISA were used for further analyses.

Table 1. FB₁ production by *F. verticillioides* isolates originating from wheat and maize grains

MRIZP Isolate Code	Origin of isolates			FB ₁ analysis (ppm)	
	Host plant	Location	Year	HPTLC	ELISA
1280	Maize	B. Topola	2006	1200	934.80
1576	Maize	B. Topola	2007	1000	1271.30
611	Maize	Kikinda	2006	100	89.40
1583	Maize	Kikinda	2007	200	206.90
1618	Maize	Kovin	2007	1000	1277.90
1558	Maize	Novi Sad	2007	100	88.60
1594	Maize	Sombor	2007	1000	1300.60
1278	Maize	S. Mitrovica	2006	200	210.53
1624	Maize	S. Mitrovica	2007	400	450.41
1411	Maize	Kraljevo	2006	200	129.00
1654	Maize	Kraljevo	2007	1000	1143.00
1651	Maize	Loznica	2007	100	88.87
1320	Maize	Niš	2006	1000	1141.60
1634	Maize	Niš	2007	1000	1141.60
1379	Maize	Šabac	2006	200	197.20
1612	Maize	Šabac	2007	400	378.40
Average				568.75	628.13
881	Wheat	Novi Sad	2005	600	659.00
675	Wheat	Sombor	2005	600	578.90
894	Wheat	S. Mitrovica	2005	400	456.70
1006	Wheat	S. Mitrovica	2006	800	728.20
1221	Wheat	Kraljevo	2006	800	882.00
804	Wheat	Šabac	2005	800	873.50
Average				666.75	696.38
Total average				617.71	662.25

Based on the ELISA test all 22 investigated isolates of *F. verticillioides* had the ability to produce toxin FB₁, but in different concentrations (Table 1). In general, the FB₁ production by this fungus varied from 88.60 ppm (MRIZP 1558) to 1,300.6 ppm (MRIZP 1594) and averaged to 662.26 ppm. Isolates of *F. verticillioides* originating from maize in 2007 were recorded in five out of the ten sites and produced more FB₁ toxin than isolates from the same location in 2006. Exceptionally, isolates originating from the location of Niš produced the same amount of this mycotoxin (1141.60 ppm) during the period of two years.

F. verticillioides isolates tested in this study had an average or high FB₁ production potential. Moreover, the highest concentration obtained in this study was lower than highest concentrations obtained in other studies, such as 4,500 ppm (STĚPIEŇ *et al.*, 2011), 4,000 ppm (STANKOVIĆ *et al.*, 2008b) and 3,800 ppm (LOGRIECO *et al.*, 2003). All produced FB₁ concentrations in this study were in the range of FB₁ production isolates tested by all above mentioned authors, and they were the closest to the FB₁ production potential of isolates originating from maize investigated in Croatia (CVETNIĆ *et al.*, 2005). The similarity between fungal species presence in maize in Serbia and Croatia, and the similarity of their mycotoxin production abilities can be explained by the influence of the similar agroecological conditions in these two neighbouring countries. Isolates with the lowest FB₁ production in this study were within the range of FB₁ concentrations reported by PLATTNER *et al.* (1990).

According to the Mann-Whitnev U test, there were no significant differences between the FB₁ production mean values of the maize and wheat originating isolates of *F. verticillioides* (628.13 ppm and 696.38 ppm, respectively). However, intraspecies variability of FB₁ production potential by *F. verticillioides* was more expressed in maize than in wheat originating isolates (Figure 1). This fact was also confirmed by C_v values (variability coefficients), which were 81.5% and 23.9% for *F. verticillioides* isolates originating from maize and wheat, respectively. DESJARDINS *et al.* (1992) also concluded that the coefficient of variation for fumonisin production on maize kernels by *F. verticillioides* isolates was high.

Isolates of *F. proliferatum* produced FB₁ toxin in a range of 2.3 ppm (MRIZP 1322) to 1,246.00 ppm (MRIZP 1345) on the average 185.20 ppm (Table 2). In four cases, *F. proliferatum* isolates originating from maize biosynthesised a higher amount of the toxin in 2006 than the isolates originating from the same location in 2007.

F. proliferatum isolates tested in this research predominantly had the average FB₁ production potential, and a maximally produced concentration was lower than the highest concentrations reported by other authors - 1,670 ppm (NELSON *et al.*, 1993), 2,250 ppm (LOGRIECO *et al.*, 1995), 2,500 ppm (LOGRIECO *et al.*, 2003), 1,820 – 2,419 ppm (STĚPIEŇ *et al.*, 2011), but within the range of FB₁ concentrations produced by isolates tested by all above mentioned authors.




Figure 1. The *F. verticillioides* isolates variability in the FB₁ production according to the host plant origin

There were no significant differences between the FB₁ production mean values of the maize and wheat originating isolates of *F. proliferatum* based on the Mann-Whitney U test (212.32 ppm and 158.07 ppm, respectively). Nevertheless, greater intraspecies variability in the FB₁ production was found between isolates of *F. proliferatum* originating from maize than those from wheat (Figure 2). These results are consistent with the C_v values, which amounted to 153.5% and 75.1% for *F. proliferatum* isolates originating from maize and wheat, respectively.

According to the Mann-Whitney U test, there were no significant differences between FB₁ production mean values of maize and wheat originating isolates of both species, but higher variability coefficients values of maize originating isolates indicate that maize is a more suitable host plant for expression of diversity in the FB₁ production than wheat. In Brazil, researchers also detected maize as a more favourable host-plant during the analysing strains isolated from maize and sorghum in relation to their production of fumonisins (DA SILVA *et al.*, 2006). In their research, maize originating strains demonstrated high levels of fumonisins, whereas the levels for sorghum originating strains were found to be low.

Table 2. *FB₁* production by *F. proliferatum* isolates originating from wheat and maize grains

MRIZP Isolate Code	Origin of isolates			FB ₁ analysis (ppm)	
	Host plant	Localition	Year	HPTLC	ELISA
1574	Maize	B. Topola	2007	100	109.90
1584	Maize	Kikinda	2007	200	205.80
1616	Maize	Kovin	2007	100	138.5
1345	Maize	Novi Sad	2006	1000	1246.00
1555	Maize	Novi Sad	2007	200	141.99
1275	Maize	Sombor	2006	200	111.20
1598	Maize	Sombor	2007	100	109.90
1653	Maize	Kraljevo	2007	trace	67.80
1248	Maize	Loznica	2006	200	219.50
1641	Maize	Loznica	2007	100	128.10
1322	Maize	Niš	2006	trace	2.37
1374	Maize	Šabac	2006	200	128.50
1609	Maize	Šabac	2007	100	76.80
Average				200.00	206.64
1432	Wheat	B. Topola	2006	trace	20.01
721	Wheat	Kikinda	2005	200	118.70
1336	Wheat	Kovin	2006	100	98.30
847	Wheat	Šabac	2005	200	260.20
674	Wheat	Sombor	2005	200	100.20
899	Wheat	S. Mitrovica	2005	200	136.31
1340	Wheat	Loznica	2006	400	372.80
Average				216.67	158.07
Total average				208.34	185.20

Table 3 shows the frequency of *F. verticillioides* and *F. proliferatum* isolates depending on the host plant origin and the level of FB₁ toxin production. The largest number of *F. verticillioides* isolates originating from the maize produced medium (40.9% of isolates) to high (31.8% of isolates) levels of FB₁. The largest number of *F. proliferatum* isolates originating from both maize (55.0%) and wheat (30.0%) produced a medium level of the FB₁ toxin. No isolate of *F. proliferatum* originating from wheat produced high levels of toxin.




Figure 2. The *F. proliferatum* isolates variability in the FB₁ production according to the host plant origin

Table 3. Categorisation of tested isolates according to the origin and the level of the FB₁ production

Host Plant	Total no. of isolates per year			FB ₁ production (ppm)*					
				No of isolates			% of isolates		
	2	0	0	Low (<50)	Medium (50-500)	High (>500)	Low (<50)	Medium (50-500)	High (>500)
	2	20	06						
	5		7						
<i>F. verticillioides</i> isolates									
Wheat	4	2	-	0	1	5	0.0	4.5	22.7
Maize	-	6	10	0	9	7	0.0	40.9	31.8
<i>F. proliferatum</i> isolates									
Wheat	4	3	-	1	6	0	5.0	30.0	0.0
Maize	-	5	8	1	11	1	5.0	55.0	5.0

*Tested isolates were divided into three groups according to Nelson *et al.* (1993)

Assessment of interspecies variability accordingly the FB₁ production potential in this study indicated that in general *F. verticillioides* isolates had a greater potential for the FB₁ production than *F. proliferatum* isolates, which was also reported by other authors (ACUÑA *et al.*, 2005; LOGRIECO *et al.*, 2003).

Both *F. verticillioides* and *F. proliferatum* isolates had a higher frequency in maize than in wheat kernels. Previous studies performed by LOGRIECO *et al.* (2003) indicated a greater variability of *Fusarium* species isolated from wheat than those isolated from maize kernels, which is in agreement with studies performed in Serbia (TANČIĆ, 2009; LEVIĆ *et al.*, 2008; KRNJAJA *et al.*, 2008; LEVIĆ *et al.*, 2004). The diversity and incidence of *Fusarium* species can be influenced not only by abiotic environmental factors but also by a host-plant. It could be considered that the presence of *F. verticillioides* and *F. proliferatum* during two years of the *Fusarium* diversity analysis on wheat kernels was a result of the environmental conditions × host plants interaction.

In this study, the FB₁ potential production of two species was analysed under optimal conditions, which may not mean that these isolates would achieve the maximum of their genetic potential under the environmental conditions. DESJARDINS *et al.* (1992) started genetic examination of the fumonisin biosynthetic pathway and studied the heritability of fumonisin production in mating population A (*F. verticillioides*), and their results indicated that variation in fumonisin levels was largely due to analytical or environmental variables. Also, these authors proved that the group of seedlings treated with fumonisin-producing progeny was significantly different from control seedlings and from the group of seedlings treated with fumonisin-nonproducing progeny in seedling emergence and in shoot length. This indicated that fumonisin plays a role in virulence but also indicated that fumonisin production is not necessary or sufficient for virulence on maize seedlings. Furthermore, if we consider the fact that the isolate MRIZP 1594 in this research had a potential to produce 1,300.6 ppm FB₁, the potential risk exists. The development of a molecular genetic analysis and genetic manipulation is a potentially powerful technique for identifying the fumonisin biosynthetic pathway and for determining the relationship between fumonisin production and the ability to cause animal and plant diseases.

CONCLUSION

Obtained results indicate that the origin of the isolates of *F. verticillioides* and *F. proliferatum* affect their potential for FB₁ biosynthesis. Namely, intraspecies variability for the mycotoxins production was greater in isolates originating from maize grain in comparison to isolates of these species from wheat grain. Interspecies variability according to the FB₁ production potential indicated that *F. verticillioides* isolates had a greater potential for the FB₁ production than *F. proliferatum* isolates. A year of the strains isolation had a distinct effect on the toxin production by *F. verticillioides* and *F. proliferatum* isolates.

ACKNOWLEDGEMENTS

This paper was realized as a part of projects TR 31023, financed by the Ministry of Education and Science of the Republic of Serbia.

Received October 10th, 2011

Accepted February 23rd, 2012

REFERENCES

- ACUÑA, A. M.C. LOZANO, M.C. DE GARCÍA, G.J. DIAZ (2005): Prevalence of *Fusarium* species of the *Liseola* section on selected Colombian animal feedstuffs and their ability to produce fumonisins. *Mycopathologia*, 160, 63-66.
- BATATINHA M.J.M., M.M.S. SIMAS, M.B. BOTURA, T.C. BITENCOURT, T.A. REIS, B. CORREA (2007): Fumonisin in brewers grain (barley) used as dairy cattle feed in State of Bahia, Brazil. *Food Control*, 18 (5): 608-612.
- BIRZELE, B., A. MEIER, H. HINDORF, J. KRÄMER, H.W. DEHNE (2002): Epidemiology of *Fusarium* infection and deoxynivalenol content in winter wheat in the Rhineland, Germany. A. LOGRIECO, J. A. BAILEY, L. CORAZZA, B. M. COOKE, eds., *Mycotoxins in Plant Disease*, Kluwer Academic Publishers, pp. 667 – 673.
- BOČAROV-STANČIĆ, A., Z. MAŠIĆ, M. ŠKRINJAR (2000): *Fusarium* species and fusariotoxins natural occurrence in cereals and commercial mixed feed originated from Jugoslavia. Proceedings of the 12th Seminar of Serbian veterinary, 78-84.
- BOTTALICO, A., G. PERRONE (2002): Toxigenic *Fusarium* species and mycotoxins associated with head blight in small-grain cereals in Europe. A. LOGRIECO, J. A. BAILEY, L. CORAZZA, B. M. COOKE eds., *Mycotoxins in Plant Disease*, Kluwer Academic Publishers, pp. 611 – 624.
- BRITZ, H., E.T. STEENKAMP, T.A. COUTHINO, B.D. WINGFIELD, W.F.O. MARASAS, M.J. WINGFIELD (2002): Two new species of *Fusarium* section *Liseola* associated with mango malformation. *Mycologia*, 94 (4), 722-730.
- CVETNIĆ, Z., S. PEPELJNIAK, M. ŠEGVIĆ (2005): Toxigenic potential of *Fusarium* species isolated from non-harvested maize. *Archives of Industrial Hygiene and Toxicology*, 56, 275-280.
- DA SILVA, V.N., F.M. FERNANDES, A. CIRTEZ, D.H. RIBEIRO, A.P. DE ALMEIDA, R.H. HASSEGAWA, B. CORREA (2006): Characterization and genetic variability of *Fusarium verticillioides* strains isolated from maize and sorghum in Brazil based on fumonisins production, microsatellites, mating type locus, and mating crosses. *Canadian Journal of Microbiology* 52 (8): 798-804.
- DESJARDINS, A.E. (2006): *Fusarium* mycotoxins: chemistry, genetics, and biology. APS Press, St. Paul, MN
- DESJARDINS, A.E., H.K. MANANDHAR, R.D. PLATTNER, G.G. MANANDAR, S.M. POLING, C.M. MARAGOS (2000): *Fusarium* species from Nepalese rice and production of mycotoxins and gibberellic acid by selected species. *Applied and Environmental Microbiology*, 66, 1020-1025.
- DESJARDINS, A.E., R.D. PLATTNER, D.D. SHACKELFORD, J.F. LESLIE, P.E. NELSON (1992): Heritability of Fumonisin B1 Production in *Gibberella fujikuroi* Mating Population A. *Applied and Environmental Microbiology*, 58 (9), 2799-2805.
- DILAS, S., B. ABRAMOVIĆ, Z. MAŠIĆ (2003): Fumonisin distribution and assessment methods. Proceedings of the 15th Serbian Veterinary Seminar, 110-118.

- FANDOHAN, P., K. HELL, W.F.O. MARASAS, M.J. WINGFIELD (2003): Infection of maize by *Fusarium* species and contamination with fumonisins in Africa. *African Journal of Biotechnology*, 2 (12), 570-579.
- FURLONG, E.B., L.M.V. SOARES, C.C. LASCA, E.Y. KOHARA (2005): Mycotoxins and Fungi in Wheat Harvested During 1990 in Test Plots in the State of São Paulo, Brazil. *Mycopathologia*, 131 (3), 185-190.
- HOLCOMB, M., J.B. SUTHERLAND, M.P. CHIARELLI, W.A. KORFMACHER, H.C. THOMPSON, J.O. JR. LAY, L.Y. HANKINS, C.E. CERNIGLIA (1993): HPLC and FAB mass spectrometry analysis of fumonisins B₁ and B₂ produced by *Fusarium moniliforme* on food substrates. *Journal of Agricultural and Food Chemistry*, 41, 357-360.
- KRNJAJA, V., J. LEVIĆ, S. STANKOVIĆ (2008): Pathogenic fungi on wheat grain in Serbia. *Journal of Plant Pathology*, 90 (3), 58.
- LE BARS, J., P. LE BARS, J. DUPUY, H. BOUDRA (1994): Biotic and abiotic factors in fumonisin B₁ production and stability. *Journal of AOAC International*, 77, 517-521.
- LESLIE, J.F., A.B. SUMMERELL (2006): *The Fusarium Laboratory Manual*. 1st Ed. Blackeell Publishing, Ames, Iowa.
- LEVIĆ, J., D. IVANOVIĆ, T. PETROVIĆ, S. STOJKOV (2001): Ambient conditions impact on fusariosis appearance on maize and mycotoxin biosynthesis, R. SPASIĆ, ed., Proceedings of the 5th Yugoslavian Plant Protection Seminar, Zlatibor, pp. 46-47.
- LEVIĆ, J., S. STANKOVIĆ, A. BOČAROV-STANČIĆ, M. ŠKRINJAR, Z. MAŠIĆ (2004): The Overview on Toxigenic Fungi and Mycotoxins in Serbia and Montenegro, A. LOGRIECO, A. VISCONTI, eds., *An Overview on Toxigenic Fungi and Mycotoxins in Europe*, Academic Publishers, London, pp. 201-218.
- LEVIĆ, J., S. STANKOVIĆ, V. KRNJAJA (2008): Harmful microorganisms in stored wheat, P. KLJAJIĆ, ed., Vizartis, Belgrade, pp. 39-66.
- LOGRIECO, A., A. BOTTALICO, G. MULÉ, A. MORETTI, G. PERRONE (2003): Epidemiology of toxigenic fungi and their associated mycotoxins for some Mediterranean crops. *European Journal of Plant Pathology*, 109, 645-667.
- LOGRIECO, A., A. MORETTI, A. RITIENI, A. BOTTALICO, P. CORDA (1995): Occurrence and toxigenicity of *F. proliferatum* from preharvest maize ear rot and associated mycotoxins in Italy. *Plant Disease*, 79, 727-731.
- LOGRIECO, A., G. MULE, A. MORETTI, A. BOTTALICO (2002): Toxigenic *Fusarium* species and mycotoxins associated with maize ear rot in Europe, A. LOGRIECO, J.A. BAILEY, L. CORAZZA, B.M. COOKE, eds., *Mycotoxins in Plant Disease*, Kluwer Academic Publishers, pp. 597 – 609.
- MARASAS, W.F.O., J.P. RHEEDER, A. LOGRIECO, P.S. VAN WYK, J.H. JUBA (1998): *Fusarium nelsonii* and *F. musarum*: two new species in Section *Arthrosporiella* related to *F. camptoceras*. *Mycologia*, 90 (3), 505-513.
- NELSON, E.P., D.R. PLATTNER, D.D. SCHACKELFORD, E.A. DESJARDINS (1991): Production of fumonisins by *Fusarium moniliforme* strains from different substrates and geographic areas. *Applied Environmental Microbiology*, 58, 984-89.
- NELSON, E.P., E.A. DESJARDINS, D.R. PLATTNER (1993): Fumonisins, mycotoxins produced by *Fusarium* species: biology, chemistry and significance. *Annual Review of Phytopathology*, 31, 233-252.

-
- PLATTNER, R.D., W.P. NORRED, C.W. BACON, A.V. KENNETH, R. PETERSON, D.D. SHACKELFORD, D. WEISLEDER (1990): A method of detection of fumonisins in corn samples associated with field cases of equine leucoencephalomalacia. *Mycologia*, 82 (6), 698-702.
- STANKOVIĆ, S., J. LEVIĆ, T. PETROVIĆ, V. KRNJAJA (2008a): Mycotoxin production by *Fusarium proliferatum* and *F. verticillioides* isolated from hops in Serbia. *Journal of Plant Pathology*, 90 (3), 58.
- STANKOVIĆ, S., J. LEVIĆ, S. TANČIĆ, V. KRNJAJA (2008b): Fumonisin B1 contamination of wheat kernels in Serbia. *Book of abstracts from IX Plant Protection Congress at Zlatibor*, 56-57.
- STANKOVIĆ, S., J. LEVIĆ, T. PETROVIĆ, V. KRNJAJA (2008c): Toxicological profile of *F. proliferatum* isolated from maize seed, root and stalk. *Cereal Research Communications*, 26 (Suppl. B), 397-398.
- STANKOVIĆ, S., J. LEVIĆ, V. KRNJAJA (2011): Fumonisin B1 in maize, wheat and barley grain in Serbia. *Biotech in Animal Husbandry* 27 (3): 631-641.
- STĘPIEŃ, Ł., G. KOCZYK, A. WAŚKIEWICZ (2011): Genetic and phenotypic variation of *Fusarium proliferatum* isolates from different host species. *J. Appl. Genet*, 52(4): 487-496.
- TANČIĆ, S. (2009): Variability of toxigenic *Fusarium* species and fusariotoxins under different agroecological maize and wheat growing conditions. Ph.D. Thesis. University of Belgrade, Serbia.
- VISCONTI, A., M.B. DOKO (1994): Survey of fumonisin production by *Fusarium* isolated from cereals in Europe. *Journal of AOAC International*, 77, 546-550.

**DIVERZITET *FUSARIUM VERTICILLIOIDES* I *F. PROLIFERATUM*
IZOLATA U POTENCIJALU SINTEZE FUMONIZINA B₁ ZAVISNO OD
POREKLA**

Sonja TANČIĆ¹, Slavica STANKOVIĆ², Jelena LEVIĆ², Vesna KRNJAJA³,
Jelena VUKOJEVIĆ⁴

¹ Laboratorija za fitopatologiju, Odeljenje za uljane culture, Institut za ratarstvo i
povrtarstvo Novi Sad, Srbija

² Laboratorija za fitopatologiju, Institut za kukuruz Zemun Polje, Srbija

³ Institut za stočarstvo, Zemun, Srbija

⁴ Katedra za algologiju, mikologiju i lihenologiju, Biološki fakultet, Univerzitet u
Beogradu, Beograd, Srbija

I z v o d

Vrste roda *Fusarium* karakteriše izuzetna intraspecijska i interspecijska varijabilnost u pogledu morfoloških, fizioloških i genetičkih svojstava. Proučavan je intraspecijski i interspecijski diverzitet izolata vrsta *Fusarium verticillioides* i *Fusarium proliferatum* u produkciji fumonizina B₁ zavisno od njihovog porekla sa zrna kukuruza i pšenice. Potencijal sinteze fumonizina B₁ kod 42 izolata je ispitivan HPTLC i ELISA metodom. Svih 22 izolata vrste *F. verticillioides* i 20 izolata vrste *F. proliferatum* su imali sposobnost stvaranja toksina fumonizina B₁. Potencijal za sintezu fumonizina je varirao od 88.60 ppm do 1,300.60 ppm kod izolata vrste *F. verticillioides* i od 2.37 ppm do 1,246.00 ppm kod *F. proliferatum* izolata. Prema Mann-Whitnev U testu nije bilo statistički značajnih razlika srednjih vrednosti sintetisanog FB₁ između *F. verticillioides* izolata poreklom sa kukuruza (628.13 ppm) i pšenice (696.38 ppm), kao ni kod *F. proliferatum* izolata poreklom sa kukuruza (212.32 ppm) i pšenice (158.07 ppm). Visoke vrednosti koeficijenta varijacije ukazuju da je varijabilnost u potencijalu sinteze fumonizina B₁ bila više izražena kod izolata poreklom sa kukuruza nego pšenice kod obe ispitivane vrste.

Primljeno 10.X. 2011.

Odobreno 23.II. 2012.