


13th
INTERNATIONAL
SYMPOSIUM

MODERN
TRENDS
IN LIVESTOCK
PRODUCTION



P R O C E E D I N G S

6 - 8 October 2021, Belgrade, Serbia

Institute for Animal Husbandry
Belgrade - Zemun, SERBIA

**13th
INTERNATIONAL
SYMPOSIUM**

**MODERN
TRENDS
IN LIVESTOCK
PRODUCTION**



6-8 October 2021, Belgrade, Serbia

PATRON

Ministry of Education, Science and Technological
Development of the Republic of Serbia

ORGANIZER

Institute for Animal Husbandry
Autoput 16, P. Box. 23,
11080, Belgrade-Zemun, Serbia
Tel: +381 11 2691 611; +381 11 2670 121; +381 11 2670 541;
Fax: + 381 11 2670 164;

PUBLISHER

Institute for Animal Husbandry, Belgrade-Zemun, Serbia
Editor-in-Chief
Čedomir Radović, PhD, Senior Research associate
Director of the Institute for Animal Husbandry, Belgrade-Zemun

Editor

Zdenka Škrbić, PhD, Principal Research Fellow
Institute for Animal Husbandry, Belgrade-Zemun

The Proceedings is printed by the Institute for Animal Husbandry,
Belgrade, 2021

biotechnology.izs@gmail.com

www.istocar.bg.ac.rs

Circulation 100 copies.

ISBN 978-86-82431-77-0



INTERNATIONAL SCIENTIFIC COMMITTEE

CHAIRMAN

Prof. Dr. **Giacomo Biagi**,
Department of Veterinary Medical Sciences, University of Bologna, Italy

SECRETARY

Dr. **Čedomir Radović**,
Institute for Animal Husbandry, Belgrade-Zemun, Serbia

MEMBERS

Dr. **Zdenka Škrbić**,
Institute for Animal Husbandry, Belgrade-Zemun, Serbia

Dr. **Veselin Petričević**,
Institute for Animal Husbandry, Belgrade-Zemun, Serbia

Prof. Dr. **Marjeta Čandek-Potokar**,
Agricultural Institute of Slovenia, Slovenia

Dr. **Giuseppe Bee**,
Agroscope Posieux, Posieux, Switzerland

Prof. Dr. **Elena Kistanova**,
Institute of Biology and Immunology of Reproduction „Kiril Bratanov“,
Sofia, Bulgaria

Prof. Dr. **Stayka Laleva**,
Agricultural Institute, Stara Zagora, Bulgaria

Prof. Dr. **Galia Zamaratskaia**,
Department of Molecular Sciences, BioCenter, Swedish University of
Agricultural Sciences, Uppsala, Sweden

Dr. **Miloš Lukić**,
Institute for Animal Husbandry, Belgrade-Zemun, Serbia

Dr. **Vlada Pantelić**,
Institute for Animal Husbandry, Belgrade-Zemun, Serbia

Prof. Dr. **Maya Ignatova**,
Institute of Animal Science, Kostinbrod, Bulgaria

Dr. **Dragana Ružić-Muslić**,
Institute for Animal Husbandry, Belgrade-Zemun, Serbia

Prof. Dr. **Marina I. Selionova**,
FSBSI-All-Russian Scientific Research Institute of Sheep and Goat Breeding,
Stavropol, Russia

Prof. Dr. **Radica Đedović**,
University of Belgrade, Faculty of Agriculture, Serbia



Prof. Dr. **Slavča Hristov**,
University of Belgrade Faculty of Agriculture, Serbia

Dr. **Dušica Ostojić Andrić**,
Institute for Animal Husbandry, Belgrade-Zemun, Serbia

Prof. Dr. **Lidija Perić**,
University of Novi Sad, Faculty of Agriculture Novi Sad, Serbia

Prof. Dr. **Wladyslav Migdal**,
Department of Animal Product Technology,
University of Agriculture in Kraków, Poland

Prof. Dr. **Danijela Kirovski**,
University of Belgrade, Faculty of Veterinary Medicine, Serbia

Prof. Dr. **Yalçın Bozkurt**,
Isparta University of Applied Science,
Department of Animal Science, Isparta, Turkey

Dr. **Snežana Mladenović Drinić**,
Maize Research Institute „Zemun Polje“, Zemun Polje, Serbia

Prof. Dr. **Randelin Dmitry Alexandrovich**,
Faculty of Biotechnology and Veterinary Medicine,
Volgograd State Agricultural University, Russia

Assoc. Prof. **Itskovich Aleksandr Yuryevich**,
Faculty of Biotechnology and Veterinary Medicine,
Volgograd State Agricultural University, Russia

Prof. Dr. **Dragan Radojković**,
University of Belgrade Faculty of Agriculture, Serbia

Prof. Dr. **Milun Petrović**,
University of Kragujevac Faculty of Agronomy, Serbia

Prof. Dr. **Dragan Glamočić**,
University of Novi Sad, Faculty of Agriculture, Serbia

Prof. Dr. **Snežana Trivunović**,
University of Novi Sad, Faculty of Agriculture, Serbia

Prof. Dr. **Predrag Perišić**,
University of Belgrade, Faculty of Agriculture, Serbia

Prof. Dr. **Vladan Bogdanović**,
University of Belgrade, Faculty of Agriculture, Serbia

Dr. **Violeta Caro Petrović**,
Institute for Animal Husbandry, Belgrade-Zemun, Serbia

Prof. Dr. **Zoran Ilić**,
University of Pristina, Faculty of Agricultural Sciences, Lešak, Serbia





Dr. Nevena Maksimović,

Institute for Animal Husbandry, Belgrade-Zemun, Serbia

Dr. Ivan Pavlović,

Scientific Veterinary Institute of Serbia, Serbia

Dr. Aleksandar Stanojković,

Institute for Animal Husbandry, Belgrade-Zemun, Serbia

Prof. Dr. Nenad Đorđević,

University of Belgrade, Faculty of Agriculture, Serbia

Dr. Zorica Bijelić,

Institute for Animal Husbandry, Belgrade-Zemun, Serbia

Dr. Violeta Mandić,

Institute for Animal Husbandry, Belgrade-Zemun, Serbia

Prof. Dr. Yusup A. Yuldashbaev,

Russian State Agrarian University,

Moscow Timiryazev Agricultural Academy,

Faculty of Animal Science and Biology, Russia

Prof. Dr. Pero Mijić,

Josip Juraj Strossmayer University of Osijek,

Faculty of Agrobiotechnical Sciences, Osijek, Croatia

Prof. Dr. Zoran Luković,

University of Zagreb Faculty of Agriculture,

Department of Animal Science and Technology, Croatia Asist.

Asist. Prof. Dubravko Škorput,

University of Zagreb Faculty of Agriculture,

Department of Animal Science and Technology, Croatia

Prof. Dr. Ivan Radović,

University of Novi Sad, Faculty of Agriculture, Serbia

Prof. Dr. Aleksandar Simić,

University of Belgrade, Faculty of Agriculture, Serbia

Prof. Dr. Nikola Pacinovski,

Ss Cyril and Methodius University in Skopje,

Institute of Animal Science, North Macedonia

Prof. Dr. Yessenbay Islamov,

Kazakh National Agrarian University, Kazakhstan

Prof. Dr. Ricmar P. Aquino,

University President, Isabela State University, Philippines

Prof. Dr. Rosa Nieto,

Departament of Physiology and Biochemistry of Animal Nutrition Estacion

Experimental del Zaidín, CSIC Armilla, Granada, Spain

Dr. Juan M. García Casco,

Departamento Mejora Genética Animal, INIA, Madrid, Spain

Dr. Slavica Stanković,

Maize Research Institute „Zemun Polje“, Zemun Polje, Serbia

Dr. Vesna S. Krnjaja,

Institute for Animal Husbandry, Belgrade-Zemun, Serbia

Prof. Dr. Rui Miguel Carracha Charneca,

Universidade de Évora, Escola de Ciências e Tecnologia,

Instituto de Ciências Agrárias e Ambientais Mediterrânicas (ICAAM), Évora,

Portugal

Dr. Jean-Louis Peyraud,

INRA, UMR PEGASE, France



ORGANIZING COMMITTEE

CHAIRMAN

Dr. Čedomir Radović,
Institute for Animal Husbandry, Belgrade-Zemun, Serbia

SECRETARY

Dr. Dragan Nikšić,
Institute for Animal Husbandry, Belgrade-Zemun, Serbia

MEMBERS

Dr. Maja Petričević,
Institute for Animal Husbandry, Belgrade-Zemun, Serbia

Dr. Marina Lazarević,
Institute for Animal Husbandry, Belgrade-Zemun, Serbia

Dr. Nikola Delić,
Institute for Animal Husbandry, Belgrade-Zemun, Serbia

Dr. Ljiljana Samolovac,
Institute for Animal Husbandry, Belgrade-Zemun, Serbia

Dr. Marija Gogić,
Institute for Animal Husbandry, Belgrade-Zemun, Serbia



SYMPOSIUM SECRETARIAT

CHAIRMAN

Dr. Dragan Nikšić,
Institute for Animal Husbandry, Belgrade-Zemun, Serbia

MEMBERS

Slavko Maletić, grad. econ.
Institute for Animal Husbandry, Belgrade-Zemun, Serbia

Olga Devečerski, grad. prof.
Institute for Animal Husbandry, Belgrade-Zemun, Serbia

Nenad Mičić, MSc,
Institute for Animal Husbandry, Belgrade-Zemun, Serbia

Bogdan Cekić, MSc,
Institute for Animal Husbandry, Belgrade-Zemun, Serbia

Miloš Marinković, MSc,
Institute for Animal Husbandry, Belgrade-Zemun, Serbia

Vladimir Živković, BSc,
Institute for Animal Husbandry, Belgrade-Zemun, Serbia

Nenad Stojiljković, BSc,
Institute for Animal Husbandry, Belgrade-Zemun, Serbia

Ivan Čosić, BSc,
Institute for Animal Husbandry, Belgrade-Zemun, Serbia

Tamara Stamenić, BSc,
Institute for Animal Husbandry, Belgrade-Zemun





Address:

Institute for Animal Husbandry,
Autoput 16, P. Box 23,
11080, Belgrade-Zemun, Serbia

Tel:

+381 11 2691 611

+381 11 2670 121

+381 11 2670 541

Fax:

+381 11 2670 164

E-mail: biotechnology.izs@gmail.com
www.istocar.bg.ac.rs

CONTENTS

INVITED PAPERS

- Marjeta Čandek-Potokar, Giuseppe Bee*
NEW TRENDS IN PUBLISHING RESEARCH AND
TRANSFERRING THE KNOWLEDGE ON ANIMAL
PRODUCTION (Slovenia-Switzerland)..... 1-10
- Vesna Gantner, Denis Kučević, Muhamed Brka*
AGRICULTURE AND ANIMAL PRODUCTION – FROM A
FOUNDER OF CIVILIZATION TO A FAILURE OR
SUSTAINABILITY (Croatia-Serbia-Bosnia and Herzegovina)..... 11-22
- Giuseppe Bee, Catherine Ollagnier*
IMPACT OF NUTRIENT SELF-SUPPLY THROUGH CHOICE
FEEDING ON GROWTH PERFORMANCE, FEEDING
BEHAVIOUR AND PROTEIN EFFICIENCY IN GROWING
FINISHING PIGS (Switzerland)..... 23-45
- Pero Mijić, Zdenko Ivkić, Tina Bobić*
RESEARCH OF PRODUCTION RESULTS IN THE
TRANSITION FROM CONVENTIONAL TO ROBOTIC
MILKING (Croatia)..... 46-55
- Ljiljana Samolovac, Slavča Hristov, Dušica Ostojić Andrić, Vlada
Pantelić, Dragan Nikšić, Dragan Stanojević, Tamara Stamenić*
ASSESSMENT OF BIOSECURITY AND WELFARE OF
CALVES REARED IN INTENSIVE HOUSING SYSTEM (Serbia). 56-72
- Marina Lazarević, Nevena Maksimović, Nenad Mičić, Miloš
Marinković, Vlada Pantelić, Dragan Nikšić, Dragan Stanojević*
THE ASSESSMENT OF BREEDING VALUE OF FIRST
CALVING HOLSTEIN - FRISIAN HEIFERS BY APPLYING
SELECTION INDEX METHODOLOGY (Serbia)..... 73-86
- Violeta Caro Petrović, Milan P. Petrović, Dragana Ružić-Muslić,
Nevena Maksimović, Marina I. Selionova, Bogdan Cekić, Ivan Ćosić*
PARITY EFFECT ON LAMBING RATE AND LAMBS BIRTH
WEIGHT (Serbia-Russia)..... 87-95
- Nikola Metodiev, Penka Moneva, Ivan Yanchev, Kostadin Kanchev*
HEMATOLOGICAL STATUS OF ILE DE FRANCE SHEEP
DEPENDING ON THEIR BODY CONDITION SCORE (Bulgaria). 96-102

<i>Nevena Maksimović, Dragana Ružić-Muslić, Violeta Caro Petrović, Bogdan Cekić, Ivan Ćosić, Nikola Delić, Marina Lazarević</i> CURRENT STATE OF GOAT BREEDING IN CENTRAL SERBIA (Serbia).....	103-116
<i>Marina I. Selionova, Tatyana V. Mamontova, Ali-Magomet M. Aybazov, Violeta Caro Petrović, Milan P. Petrović</i> QUALITY OF ABORIGENOUS KARACHAY GOAT MEAT UNDER DIFFERENT CONDITIONS (Russia-Serbia).....	117-125
<i>Nataša Tolimir, Zdenka Skrbić, Marijana Maslovarić, Miloš Lukić, Dragan Milić</i> THE IMPORTANCE OF EGGS IN THE DIET, CONSUMER PREFERENCES, THE PRODUCTION AND MARKET OF TABLE EGGS IN SERBIA (Serbia).....	126-148
<i>Radomir Savić, Vesna Davidović, Ivana Božičković</i> ASSESSMENT OF FERTILITY OF BOARS – DIFFERENT APPROACHES (Serbia).....	149-162
<i>Zoran Luković, Danijel Karolyi, Sven Menčik, Dubravko Škorput</i> EFFECT OF BIRTH WEIGHT ON SURVIVAL AND GROWTH PERFORMANCE OF PIGLETS FROM LARGE LITTERS (Croatia).....	163-172
<i>Patricia Palma-Granados, Ignacio Fernández-Fígares, Isabel Seiquer, Manuel Lachica, Luis Lara, Ana Haro, Rosa Nieto</i> PERFORMANCE, METABOLIC AND MEAT QUALITY IMPLICATIONS OF IMMUNOCASTRATION IN IBERIAN PIGS (Spain).....	173-184
<i>Galia Zamaratskaia, Andriy Getya</i> RAISING MALE PIGS AS AN ALTERNATIVE TO SURGICAL CASTRATION (Sweden-Ukraine).....	185-194
<i>Władysław Migdał, Rafał Duś, Maria Walczycka, Łukasz Migdał</i> SILESIA PIGS' FATTENERS - THE IDEA FOR HEALTHY PIG AND HIGH QUALITY PORK MEAT (Poland).....	195-207
<i>Marija Gogić, Nenad Katanić, Vladimir Živković, Nenad Stojiljković, Violeta Mandić, Maja Petričević, Radomir Savić</i> LIFE DAILY GAIN OF INDIGENOUS PIG BREEDS IN SERBIA (Serbia).....	208-216

<p><i>Nikola Delić, Dragan Nikšić, Maja Petričević, Aleksandar Stanojković, Vladimir Živković, Marina Lazarević, Nevena Maksimović</i> THE EFFECT OF PHYTOGENIC ADDITIVES ON THE DEGREE OF BACTERIAL INFECTION <i>B. HYODYSENTERIAE</i> IN WEANED PIGLETS (Serbia).....</p>	217-226
<p><i>Costanza Delsante, Carlo Pinna, Federica Sportelli, Claudio Stefanelli, Carla G. Vecchiato, Giacomo Biagi</i> ASSESSMENT OF THE EFFECTS OF EDIBLE MICROALGAE IN A CANINE GUT MODEL (Italy).....</p>	227-242
<p><i>Yalcin Bozkurt, Serkan Ozkaya, Sukran Kuleasan</i> THE EFFECTS OF PRE-SLAUGHTER RESTING PERIODS ON CARCASS AND MEAT QUALITY DETERMINED BY DIGITAL IMAGE ANALYSIS (Turkey).....</p>	243-251
<p><i>Nikola Stanišić, Milo Mujović, Slaviša Stajić, Maja Petričević, Čedomir Radović, Marija Gogić, Aleksandar Stanojković</i> TECHNO-FUNCTIONAL PROPERTIES OF THREE DIETARY FIBERS USED IN THE MEAT PROCESSING INDUSTRY (Netherlands-Serbia).....</p>	252-262
<p><i>Slaviša Stajić, Ana Kalušević</i> TECHNOLOGICAL AND SENSORY PROPERTIES OF SERBIAN TRADITIONAL MINCED MEAT PRODUCT ČEVAPI WITH IMPROVED NUTRITIONAL PROPERTIES (Serbia).....</p>	263-274
<p><i>Maja Petričević, Tamara Stamenić, Dragan Nikšić, Ljiljana Samolovac, Veselin Petričević, Marija Gogić, Violeta Mandić</i> EXAMINATION OF CERTAIN OF BEEF MEAT QUALITY TRAITS UNDER THE INFLUENCE OF FLAXSEED DIET (Serbia).....</p>	275-286
<p><i>Vesna Dragičević, Milena Simić, Milan Brankov, Milena Šenk, Vesna Krnjaja, Violeta Mandić, Branka Kresović</i> BIOFORTIFICATION AS A WAY OF NUTRIENT DENSE FEED PRODUCTION (Serbia).....</p>	287-307
<p><i>Jordan Marković, Jasmina Milenković, Snežana Anđelković</i> ALFALFA AND RED CLOVER AS A PROTEIN SOURCE FOR RUMINANTS (Serbia).....</p>	308-321

Violeta Mandić, Snežana Đorđević, Zorica Bijelić, Vesna Krnjaja, Aleksandar Simić, Marija Gogić, Maja Petričević
INOCULATION OF MAIZE WITH PGPR (Serbia)..... 322-334

Maya M. Ignatova, Nadezhda M. Sertova
MYCOTOXIN CONTAMINATION OF CEREAL GRAINS IN BULGARIA (Bulgaria)..... 335-345

ORAL PRESENTATION

Slavča Hristov, Branislav Stanković, Dušica Ostojić Andrić, Ljiljana Samolovac, Nevena Maksimović, Marko Cincović, Dimitar Nakov
INFLUENCE OF CATTLE BREEDING CONDITIONS ON REPRODUCTION, GROWTH, MILK YIELD AND MEAT AND MILK QUALITY (Serbia-North Macedonia)..... 346-362

Ivan Pavlović, Violeta Caro Petrović, Dragana Ružić Muslić, Jovan Bojkovski, Nemanja Zdravković, Renata Relić, Vukašin Stefanović
GASTROINTESTINAL HELMINTHS OF SHEEP BREED IN POMORAVSKI AND RASINA DISTRICT (Serbia)..... 363-370

Jovan Bojkovski, Marina Spinu, Mihaela Niculae, Jasna Prodanov-Radulović, Aleksandar Stanojković, Ivan Pavlović, Nemanja Zdravković, Radiša Prodanović
PRODUCTION RELATED DISEASES (TECHNOPATHIES) IN SWINE COMMERCIAL FARM (Serbia-Romania)..... 371-390

Teodora Popova, Nadezhda Palova, Jivko Nakev, Maya Ignatova
CHEMICAL COMPOSITION AND FATTY ACID PROFILE OF TWO TRADITIONAL BULGARIAN DRY-CURED MEAT PRODUCTS MADE OF EAST BALKAN PIG (Bulgaria)..... 391-404

POSTER PRESENTATION

Branislav Stanković, Slavča Hristov, Dušica Ostojić Andrić, Milošević-Stanković Ivana, Dimitar Nakov, Rašović Bojanić Mirjana
PRECISION FARMING IN IMPROVEMENT OF DAIRY CATTLE WELFARE (Serbia-North Macedonia)..... 405-421

<i>Tina Bobić, Andrea Bejteš, Pero Mijić, Vesna Gantner, Maja Gregić</i> LAMENESS DETECTION IN CATTLE USING ICT TECHNOLOGY (Croatia).....	422-433
<i>Mirna Gavran, Danko Šinka, Vesna Gantner</i> THE EFFECT OF SEASON ON THE ESTIMATED AMMONIA EMISSION OF HOLSTEIN FIRST PARITY COWS (Croatia).....	434-441
<i>Franjo Poljak, Marija Špehar, Zvonimir Steiner, Mirna Gavran, Vesna Gantner</i> THE VARIABILITY IN THE ESTIMATED PREVALENCE RISK OF METABOLIC DISORDERS (KETOSIS/ACIDOSIS) IN SIMMENTAL FIRST PARITY COWS DUE TO RECORDING SEASON (Croatia).....	442-450
<i>Ivana Jožef, Dragan Solić, Zvonimir Steiner, Vesna Gantner</i> THE EFFECT OF RECORDING SEASON ON THE ESTIMATED PREVALENCE RISK OF MASTITIS IN SIMENTAL FIRST PARITY COWS (Croatia).....	451-457
<i>Marinela Enculescu</i> INVESTIGATIONS ON HAEMATO-BIOCHEMICAL INDICATORS IN ROMANIAN BLACK AND SPOTTED DAIRY COWS WITH RETAINED PLACENTA – PRELIMINARY RESULTS (Romania).....	458-467
<i>Zhivko I. Ducheve</i> ASSESSMENT OF THE RISK STATUS OF FOUR LOCAL BULGARIAN BREEDS BASED ON THEIR GEOGRAPHIC DISTRIBUTION (Bulgaria).....	468-476
<i>Tamara Papović, Denis Kučević, Miroslav Plavšić, Snežana Trivunović, Ksenija Čobanović</i> PREDICTION OF AMMONIUM EMISSION FROM DAIRY CATTLE BASED ON MILK UREA NITROGEN USING THE PRECISION FARMING METHODOLOGY (Serbia).....	477-486
<i>Dragana Ružić-Muslić, Bogdan Cekić, Ivan Ćosić, Ivan Pavlović, Nevena Maksimović, Violeta Caro Petrović, Zorica Bijelić</i> HEMATOLOGICAL AND BIOCHEMICAL BLOOD PARAMETERS OF PIROT PRAMENKA - ENDANGERED SHEEP POPULATION (Serbia).....	487-499

<i>Bogdan Cekić, Dragana Ružić Muslić, Nevena Maksimović, Violeta Caro Petrović, Ivan Ćosić, Tamara Stamenić, Madlena Andreeva</i> IMPORTANCE, PRODUCTIVITY AND POTENTIALS OF LOCAL SERBIAN SHEEP BREEDS (Serbia-Bulgaria).....	500-508
<i>Rossen Stefanov, Madlena Andreeva</i> STUDY OF THE SPERM PARAMETERS OF RAMS' EJACULATES OBTAINED IN THE BREEDING AND NON- BREEDING PERIOD (Bulgaria).....	509-515
<i>Vladimir Dosković, Snežana Bogosavljević-Bošković, Božidar Milošević, Zdenka Škrbić, Miloš Lukić, Simeon Rakonjac, Veselin Petričević</i> EFFECTS OF GENOTYPE AND PROTEASE ENZYME SUPPLEMENTATION ON THE PRIMAL CARCASS CUTS OF CHICKENS (Serbia).....	516-523
<i>Marija Pavlović, Ksenija Nešić, Aleksandra Tasić, Nikola Čobanović, Mihajlo Vićentijević, Ivan Pavlović</i> IMPACT OF INORGANIC PHOSPHORUS IN BROILERS DIET ON BONE MINERALIZATION (Serbia).....	524-531
<i>Aleksandar Pavličević, Ivan Pavlović, Nemanja Zdravković, Luis Francisco Angeli</i> Alves SUGGESTED APPROACH TO RED POULTRY MITE CONTROL IN EXTENSIVE POULTRY PRODUCTION (Serbia- Brazil).....	532-540
<i>Maja Gregić, Mirjana Baban, Pero Mijić, Vesna Gantner, Tina Bobić</i> THE CHALLENGES OF JUMPING HORSES THROUGH THE TRAINING (Croatia).....	541-551
<i>Nemanja Zdravković, Oliver Radanović, Milan Ninković, Radoslava Savić-Radovanović, Nataša Rajić Savić, Đordje Marjanović, Jovan Bojkovski</i> ACTIVITY OF SOME PLANT ESSENTIAL OILS AGAINST COMMON ISOLATES IN VETERINARY BACTERIOLOGY - A PILOT STUDY (Serbia).....	552-560
<i>Urška Tomažin, Klavdija Poklukar, Martin Škrlep, Nina Batorek Lukač, Marjeta Čandek-Potokar</i> THE EFFECT OF RYR1 GENE ON MEAT QUALITY IN AUTOCHTHONOUS BREED KRŠKOPOLJE PIG (Slovenia).....	561-571

<i>Klavdija Poklukar, Marjeta Čandek-Potokar, Milka Vrecl Fazarinc, Nina Batorek Lukač, Gregor Fazarinc, Kevin Kress, Volker Stefanski, Martin Škrlep</i> EFFECTS OF ANDROGEN DEPRIVATION ON HISTOMORPHOLOGICAL PROPERTIES OF FAT TISSUE IN PIGS (Slovenia-Germany).....	572-580
<i>Aleksandra Petrović, Dragan Radojković, Čedomir Radović, Marija Gogić, Nenad Stojiljković, Nenad Parunović, Radomir Savić</i> IN VITRO BOAR FERTILITY DURING SUMMER AND AUTUMN SEASON (Serbia).....	581-589
<i>Nenad Stojiljković, Dragan Radojković, Čedomir Radović, Marija Gogić, Vladimir Živković, Zoran Luković, Dubravko Škorput</i> VARIABILITY OF THE NUMBER OF LIVE-BORN PIGLETS UNDER THE INFLUENCE OF FEMALE GENOTYPE, YEAR OF FARROWING AND PARITY (Serbia-Croatia).....	590-597
<i>Dragan Dokić, Maja Gregić, Mirna Gavran, Vesna Gantner</i> EFFECTS OF INVESTMENTS IN CAPITAL CROP PRODUCTION - A COMPARATIVE ANALYSIS OF THE REPUBLIC OF CROATIA AND THE EUROPEAN UNION (Croatia).....	598-605
<i>Sanja Živković, Tanja Vasić</i> MYCOPOPULATION OF ALFALFA AND RED CLOVER HAY IN SERBIA (Serbia).....	606-614
<i>Ksenija Nešić, Nikola Pavlović, Marija Pavlović, Jelena Vlajković, Aleksandra Tasić, Vladimir Radosavljević, Božidar Savić</i> AN INSIGHT INTO THE MYCOTOXICOLOGICAL SITUATION – RECENT EXPERIENCE AND CLOSE PREDICTION (Serbia)....	615-521
<i>Vesna Krnjaja, Violeta Mandić, Zorica Bijelić, Slavica Stanković, Milica Nikolić, Tanja Vasić, Nikola Delić</i> FUSARIUM SPP. AND DEOXYNIVALENOL CONTAMINATION OF RYEGRASS SEEDS (Serbia).....	622-631

EFFECTS OF GENOTYPE AND PROTEASE ENZYME SUPPLEMENTATION ON THE PRIMAL CARCASS CUTS OF CHICKENS

Vladimir Dosković¹, Snežana Bogosavljević-Bošković¹, Božidar Milošević², Zdenka Škrbić³, Miloš Lukić³, Simeon Rakonjac¹, Veselin Petričević³

¹ Faculty of Agronomy, University of Kragujevac, Cara Dušana 34, 32000 Čačak, Serbia

² Faculty of Agriculture, University of Pristina, Kosovska Mitrovica-Zubin potok-Lešak, 38219 Lešak

³ Institute for Animal Husbandry, Autoput 16, P. Fah 23, 11080, Belgrade-Zemun, Serbia

Corresponding author: Vladimir Dosković, vladosko@kg.ac.rs

Abstract: This study compared weight and percent yields of primal carcass cuts of commercial fast-growing broiler - Cobb 500 and medium-growing broiler genotype - Master Gris. Three hundred Cobb 500 chickens and three hundred Master Gris chickens were used for the experiment. Feed and water were provided *ad-libitum*. Treatments were: T1 - corn-soybean meal based control diet; T2: T1 – 4.0% crude protein than in the control diet + 0.2% protease and T3: T1 – 6.0% crude protein than in the control diet + 0.3% protease. Chickens were slaughtered at 49 days of age. Medium-growing hybrid Master Gris had a lower weight of all primal carcass cuts and breast yield (as a percentage of ready-to-grill carcass weight), as well as higher thighs, drumsticks, wings and pelvis yields in comparison with the Cobb 500 broilers ($P < 0.05$), except back yield ($P > 0.05$). Also, the addition of protease to broiler feed had no significant effect on the weight and percent yields of primal carcass cuts ($P > 0.05$).

Keywords: hybrids, broilers, protease enzyme, primal carcass cuts

Introduction

In the production of chicken meat in recent decades, increasing attention is paid to minimizing the cost of broiler fattening and reducing the environmental pollution. Several factors such as genotype, diet composition, digestible nutrient content, energy to protein ratio, feed form, feed processing, environment, and disease could affect the cost of production and poultry product quality through influencing feed intake, body weight gain and feed conversion ratio (*Ahiwe et al., 2018*).

Today, technologies supplemented by exogenous enzymes (xylanase, phytase, glucanase, protease, amylase,...) in the corn-soybean based diet are most commonly used to achieve these goals. Aiming to increase the efficiency of rations, the usage of exogenous enzymes in the feeding of broilers is gaining more space and has become a great alternative, since it enhances food digestibility, minimizing the anti-nutritional effects and promoting the productivity indexes (*Hooge et al., 2010*). These enzymes can be included in the diet of livestock to improve digestibility, reduce nitrogen and phosphorus pollution, reduce diet cost, and improve the uniformity of animal growth (*Cowieison et al., 2019*).

Proteases may have an 'extra-proteinaceous' advantage for broilers reared under high-temperature conditions by improving gut health (*Cowieison et al., 2016*). Single exogenous protease, when added to the diet, effectively increased the apparent amino acid and protein digestibility in the broiler chicken's diet, even in a low protein diet (*Romero et al., 2014; Law et al., 2015; Bertechini et al., 2020; Da Silva et al., 2021,...*). Some researchers (*Sabino et al., 2004; Kamran et al., 2008;...*) have examined the effects of reducing the levels of crude protein in the diet with exogenous enzyme supplementation on carcass yields and cuts of chickens.

The objective of this study was to compare the carcass characteristics and meat quality of a medium-growing broiler - Master Gris with a fast-growing commercial Cobb 500 broiler, both grown in identical housing conditions and fed the same diet (a diet based on maize and soybean meal, with or without protease enzymes).

Materials and Methods

A total of 300 Cobb 500 broiler chickens (1 day old, 44.73 g weight) and 300 Master Gris chickens (1 day old, 43.03 g weight) were used in the experiment. Diets were provided in phases: starter (day 1-21), grower (day 22-35) and finisher (day 35-49). Respective diets (mash feed) and water were provided *ad libitum* from day 1 to 49.

Diet treatments were T1: control diet (based on maize and soybean meal, starter phase 22% crude protein, grower phase 19% crude protein, finisher phase 17% crude protein), T2: T1 – 4.0% crude protein than in the control diet + 0.2% protease and T3: T1 – 6.0% crude protein than in the control diet + 0.3% protease. The enzyme preparation used in this study was a commercial product (Ronozyme ProAct®, produced by DSM, The Netherlands) produced through submerged fermentation of *Bacillus licheniformis*.

At the end of the experimental period (day 49), a total of 20 birds (10 male and 10 female) were randomly selected from each treatment. Chickens were

weighed and slaughtered and then the ready-to-grill carcass weight. After that, the carcasses are cut into primal cuts - breast, drumsticks, thighs, wings, back, pelvis and abdominal fat. Primal cuts is the measured weights and the proportions of primal cuts in relation to the ready-to-grill carcass weight were calculated.

Statistical analysis data were subjected to two-way factorial ANOVA (hybrids, diet treatments). The significance of differences between means was determined by the LSD test and differences were considered significant at $P < 0.05$. Statistical analyses were performed using the Stat Soft Inc Statistica For Windows, Version 7.0., 2006.

Results and Discussion

Table 1 presents weight of primal carcass cuts - breast, drumsticks, thighs, wings, back and pelvis in broilers two hybrids.

Table 1. Effect of hybrids and diet on weights of primal carcass cuts across treatments, g

Hybrid	Diet		Breast	Drumsticks	Thighs	Wings	Back	Pelvis
Cobb 500	T1	\bar{x}	798.68 ^a	326.45 ^a	376.97 ^a	261.88 ^a	283.53 ^a	246.21 ^a
		Sd	80.28	43.93	43.89	27.22	33.25	31.64
	T2	\bar{x}	791.36 ^a	312.57 ^a	372.29 ^a	252.15 ^a	274.78 ^a	244.96 ^a
		Sd	91.38	34.71	37.13	27.11	51.96	30.86
	T3	\bar{x}	764.78 ^a	307.96 ^a	361.19 ^a	253.83 ^a	279.95 ^a	252.22 ^a
		Sd	88.92	40.65	39.29	30.17	34.80	41.06
Master Gris	T1	\bar{x}	524.51 ^b	277.48 ^b	303.78 ^b	221.74 ^b	222.64 ^b	211.47 ^b
		Sd	38.66	28.44	31.98	19.49	27.09	16.44
	T2	\bar{x}	509.90 ^b	264.66 ^b	293.25 ^b	219.16 ^b	209.10 ^b	200.89 ^b
		Sd	48.88	32.09	26.78	20.41	19.00	18.15
	T3	\bar{x}	515.70 ^b	265.04 ^b	297.08 ^b	219.18 ^b	216.55 ^b	205.55 ^b
		Sd	50.58	31.26	28.49	23.41	24.80	22.06
p-value								
Source of variation								
Hybrid			0.000	0.000	0.000	0.000	0.000	0.000
Diet			0.394	0.113	0.347	0.491	0.332	0.552
Hybrid x Diet			0.554	0.921	0.633	0.799	0.950	0.608

\bar{X} -Average, Sd - Standard deviation

a-b Means followed by different superscript letters within columns differ significantly ($P < 0.05$)

The results presented in Table 1 showed a significant effect of genotype, ie hybrid on the weight of primal carcass cuts in chickens ($P < 0.05$). Average weight of primary cuts were higher in broilers Cobb 500, which was expected, according to a higher body mass chickens of hybrids at the end of the experiment. Also, the results

of the experiment show that lowering crude proteins levels, with the addition of protease enzymes, did not have a negative effect on examined carcass characteristics ($P>0.05$). The breast weight of fast-growing broilers (784.94 g) was 1.52 time higher compared to the breast weight (516.70 g) of medium-growth hybrids. *Aksoy et al. (2010)* also reported fast-growing genotypes showed higher breast, legs, wings, back and neck weight than medium growing birds ($P<0.001$). Commercial fast-growing chickens had higher live weight, carcass, breast and thigh-drumstick weights ($P<0.05$) compared to medium and slow-growing chickens (*Cömert et al., 2016*).

Data on percent yields of primal carcass cuts (breast, drumsticks, thighs, wings, back and pelvis) the factors analyzed (hybrids, dietary treatments) are given in Table 2.

Table 2. Effect of hybrids and diet on percentage yield of primal carcass cuts of broilers, %

Hybrid	Diet		Breast	Drumsticks	Thighs	Wings	Back	Pelvis
Cobb 500	T1	\bar{x}	33.60 ^a	13.69 ^b	15.83 ^b	11.02 ^b	11.92	10.34 ^c
		Sd	1.59	0.87	0.68	0.53	0.80	0.70
	T2	\bar{x}	33.93 ^a	13.38 ^b	15.96 ^b	10.79 ^b	11.68	10.47 ^{bc}
		Sd	3.59	0.90	1.17	0.65	1.46	0.76
	T3	\bar{x}	33.24 ^a	13.34 ^b	15.47 ^b	11.02 ^b	12.15	10.91 ^b
		Sd	2.29	0.67	1.19	0.62	0.66	0.96
Master Gris	T1	\bar{x}	28.62 ^b	15.10 ^a	16.53 ^a	12.08 ^a	12.11	11.54 ^a
		Sd	1.33	0.56	0.75	0.44	0.88	0.56
	T2	\bar{x}	28.90 ^b	14.97 ^a	16.62 ^a	12.42 ^a	11.86	11.40 ^a
		Sd	1.20	0.79	0.45	0.41	0.60	0.68
	T3	\bar{x}	28.78 ^b	14.77 ^a	16.58 ^a	12.22 ^a	12.08	11.47 ^a
		Sd	1.25	0.848	0.58	0.49	0.81	0.58
p-value								
Source of variation								
Hybrid			0.000	0.000	0.000	0.000	0.549	0.000
Diet			0.659	0.148	0.388	0.819	0.228	0.195
Hybrid x Diet			0.785	0.845	0.431	0.099	0.782	0.134

\bar{X} - Average, Sd - Standard deviation

a-b Means followed by different superscript letters within columns differ significantly ($P<0.05$)

The results presented in Table 1 showed that chickens of the fast-growing hybrid had a higher percent yields of breasts, and a lower percent yields of drumsticks, thighs, wings and pelvis compared to the medium-growing genotype ($P<0.05$), with a similar percent yields of backs ($P>0.05$). Nutritive applied treatments did not affect the percentage yield of the primary carcass cuts, in

addition to the percent yield of the pelvis in the genotype Cobb 500 (significant differences were observed between the groups T1 and T3, $P < 0.05$).

Relative weights of parts to cold carcass weight fast-growing genotypes showed a higher average for breast only whereas medium-growing genotypes had higher values for other parts (Aksoy *et al.*, 2010). A similar difference in carcass traits between fast- and medium-growing genotypes was also found Santos *et al.* (2004), Fanatico *et al.* (2008) and Dal Bosco *et al.* (2014). Higher relative breast weight in fast-growing broiler in comparison with slow-growing broiler were also cited by Nielsen *et al.* (2003), Fanatico *et al.* (2005) and Jaspal *et al.* (2020). Cömert *et al.* (2016) published that the effect of the genotype was important for the breast and thigh-drumstick yields ($P < 0.05$), fast-growing hybrids had higher breast yield, whereas medium slow-growing chickens had higher thigh-drumstick yield ($P < 0.05$).

Conclusion

Based on the findings of the present study, it may be concluded that the differences in weight and percent yields of primal carcass cuts between the two hybrids of different intensity of growth were significant, except for the share of the back. Namely, the weights of all primal carcass cuts and breast yield were greater in fast-growth broilers, while the percent yield of drumsticks, thighs, wings and pelvis were greater in medium growth broilers. Also, present results showed that it was possible to reduce crude protein level without any negative effect on the weight and percent yields of primal carcass cuts of commercial fast-growing broiler (Cobb 500) and medium-growing broiler genotype (Master Gris) by adding protease to the diet of broiler chickens.

Uticaj genotipa i dodatka enzima proteaze na karakteristike trupa pilića

Vladimir Dosković, Snežana Bogosavljević-Bošković, Božidar Milošević, Zdenka Škrbić, Miloš Lukić, Simeon Rakonjac, Veselin Petričević

Rezime

U radu je upoređena masa i udeo osnovnih delova trupa komercijalnog brzorastućeg hibrida pilića - Cobb 500 i srednje rastućeg linijskog hibrida - Master Gris. U ogledu je bilo tri stotine pilića Cobb 500 i tri stotine pilića Master Gris. Ishrana i napajanje su bili po volji. Tretmani ishrane su bili: T1 - kontrolni obrok

na bazi kukuruza i sojine sačme; T2: T1 – 4,0% sirovih proteina u odnosu na kontrolni obrok + 0,2% proteaze i T3: T1 – 6,0% sirovih proteina u odnosu na kontrolni obrok + 0,3% proteaze. Klanje pilića obavljeno je u uzrastu od 49 dana. Medium-growing hibrid Master Gris je imao manju masu svih osnovnih delova trupa i udeo grudi (u odnosu na masu trupa “spremno za roštilj”), kao i veći udeo bataka, karabataka, krila i karlice u poređenju sa pilićima Cobb 500 ($P < 0.05$), izuzev udela leđa ($P > 0.05$). Takođe, dodatak proteaze u hranu za piliće nije imao značajan uticaj na masu i udeo osnovnih delova trupa ($P > 0.05$).

Ključne reči: hibridi, brojleri, enzim proteaza, osnovni delovi trupa.

Acknowledgment

The research was financed by the Ministry of Education, Science and Technological Development, Republic of Serbia, project ref. number 451-03-9/2021-14 and 451-03-9/2021-14/200022.

References

- AHIWE E.U., OMEDE A.A., ABDALLH M.B., IJI P.A. (2018): Managing dietary energy intake by broiler chickens to reduce production costs and improve product quality. *Animal Husbandry and Nutrition*, Edited by Banu Yücel and Turgay Taşkin, IntechOpen Limited, London, United Kingdom, doi: 10.5772/intechopen.76972, 115-145.
- AKSOY T., NARINE D., CUREK D.I., ONENC A., YAPICI N. (2010): Comparison of fast and medium- growing broiler genotypes raised in-doors: growth performance, slaughter results and carcass parts. *Journal of Animal and Veterinary Advances*, 9, 1485-1490.
- BERTECHINI G.A., DE CARVALHO C.C.J., CARVALHO C.A., DALOLIO S.F., SORBARA O.B.J. (2020): Amino acid digestibility coefficient values of animal protein meals with dietary protease for broiler chickens. *Transl. Anim. Sci.*, 4:1-11.
- COWIESON A., ZAEFARIAN F., KNAP I., RAVINDRAN V.R. (2016): Interactive effects of dietary protein concentration, a mono-component exogenous protease and ascorbic acid on broiler performance, nutritional status and gut health. *Anim. Produ. Sci.*, 57, 1058-68.
- COWIESON A.J., SMITH A., SORBARA J.O.B., PAPPENBERGER G., OLUKOSI O.A. (2019): Efficacy of a mono-component exogenous protease in the presence of a high concentration of exogenous phytase on growth performance of broiler chickens. *J. Appl. Poult. Res.*, 28, 638-646.

- CÖMERT M., ŞAYAN Y., KIRKPINAR F., BAYRAKTAR Ö.H., MERT S. (2016): Comparison of carcass characteristics, meat quality, and blood parameters of slow and fast grown female broiler chickens raised in organic or conventional production system. *Asian-Australas J. Anim. Sci.*, 29, 7, 987-997.
- DA SILVA J.M.S., DE OLIVEIRA N.R., GOUVEIA A.B.V.S., VIEIRA R.A., DOS SANTOS R.O.F., MINAFRA C.S., DOS SANTOS F.R. (2021): Effect of protease supplementation on the digestibility of amino acids in animal-origin meals for broiler diets. *Czech J. Anim. Sci.*, 66, 29-37.
- DAL BOSCO A., MUGNAI C., GUARINO AMATO M., PIOTTOLI L., CARTONI A., CASTELLINI C. (2014): Effect of slaughtering age in different commercial chicken genotypes reared according to the organic system: 1. welfare, carcass and meat traits. *Italian Journal of Animal Science*, 13, 2, 467-472.
- FANATICO A.C., CAVITT L.C., PILLAI P.B., EMMERT J.L., OWENS C.M. (2005): Evaluation of slower-growing broiler genotypes grown with and without outdoor access: Meat quality. *Poult. Sci.*, 84, 1785-1790.
- FANATICO A.C., PILLAI P.B., HESTER P.Y., FALCONE C., MENCH J., OWENS C.M., EMMERT J.L. (2008): Performance, livability, and carcass yield of slow- and fast-growing chicken genotypes fed low-nutrient or standard diets and raised indoors or with outdoor access. *Poult. Sci.*, 87, 1012-1021.
- HOOGE D.M., PIERCE J.L., McBRIDE K.W., RIGOLIN P.J. (2010): Meta-analysis of broiler chicken trials using diets with or without allzyme SSF enzyme complex. *International Journal of Poultry Science*, 9, 9, 819-823.
- JASPAL M.H., ALI S., RAJPUT N., NAEEM M., TALPUR F.N., REHMAN I. (2020): Fatty acid profiling and comparative evaluation of carcass cut up yield, meat quality traits of Cobb Sasso, commercial broiler and native aseel chicken. *Pure Appl. Biol. (Pab)* 9, 56-65.
- KAMRAN Z., SARWAR M., NISA M., NADEEM M.A., MAHMOOD M.E. (2008): Effect of low-protein diets having constant energy-to-protein ratio on performance and carcass characteristics of broiler chickens from one to thirty-five days of age. *Poult. Sci.* 87, 468-474.
- LAW F.L., ZULKIFLI I., SOLEIMANI A.F., HOSSAIN M., LIANG J.B. (2015): Nutrient digestibility of broiler chickens fed on a low-protein diet supplemented with mono-component proteases. *Eur. Poult. Sci.*, 79, 107.
- NIELSEN B.L., THOMSEN M.G., SØRENSEN J.P., YOUNG J.F. (2003): Feed and strain effects on the use of outdoor areas by broilers. *Brit. Poult. Sci.*, 44, 2, 161-169.
- ROMERO L.F., SANDS J.S., INDRAKUMAR S.E., PLUMSTEAD P.W., DALSGAARD S., RAVINDRAN V. (2014): Contribution of protein, starch, and fat to the apparent ileal digestible energy of corn- and wheat-based broiler diets in

response to exogenous xylanase and amylase without or with protease. *Poult. Sci.* 93, 1-13.

SABINO H.F.N., SAKOMURA N.K., NEME R., FREITAS E.R. (2004): Protein levels in broiler feed during the growth phase. *Brazilian Agricultural Research* 39(5), 407-412.

SANTOS A.L., SAKOMURA N.K., FREITAS E.R., BARBOSA N.A.A., MENDONÇA M.O., CARRILHO E.N.V.M. (2004): Carcass yield and meat quality of three strains of broiler chicken. *Proceedings of 22nd World's Poultry Congress*, June 8-13, Istanbul, Turkey, pp.338.

=====
CIP - Каталогизација у публикацији
Народна библиотека Србије, Београд

636/638(082)(0.034.2)
631/635(082)(0.034.2)

**INTERNATIONAL Symposium Modern Trends in Livestock
Production (13 ; 2021 ; Beograd)**

Proceedings [Elektronski izvor] / 13th International Symposium Modern
Trends in Livestock Production, 6 -8 October 2021, Belgrade, Serbia ;
[organizer] Institute for Animal Husbandry, Belgrade - Zemun ; [editor
Zdenka Škrbić]. - Belgrade : Institute for Animal Husbandry, 2021 (Belgrade
: Institute for Animal Husbandry). - 1 USB fleš memorija ; 1 x 3 x 6 cm

Sistemski zahtevi: Nisu navedeni. - Nasl. sa naslovne strane dokumenta. -
Tiraž 100. - Bibliografija uz svaki rad.

ISBN 978-86-82431-77-0

а) Сточарство -- Зборници б) Пољопривреда -- Зборници

COBISS.SR-ID 46411785

=====



13th INTERNATIONAL SYMPOSIUM
MODERN TRENDS IN LIVESTOCK PRODUCTION

6 - 8 October 2021 - Belgrade, Serbia

P R O C E E D I N G S

INSTITUTE FOR ANIMAL HUSBANDRY

Autoput 16, P. Box 23, 11080, Belgrade - Zemun, Serbia

www.istocar.bg.ac.rs

ISBN 978-86-82431-77-0

